

ENHANCEMENT AND SUPPRESSION OF HEAT TRANSFER BY MHD TURBULENCE

A. LAZARIAN

Dept. of Astronomy, University of Wisconsin, Madison, WI53706; lazarian@astro.wisc.edu
Draft version February 5, 2008

ABSTRACT

We study of the effect of turbulence on heat transfer within magnetized plasmas for energy injection velocities both larger and smaller than the Alfvén speed. We find that in the latter regime the heat transfer is partially suppressed, while in the former regime the effects of turbulence depend on the intensity of driving. In fact, the scale l_A at which the turbulent velocity is equal the Alfvén velocity is a new important parameter. When the electron mean free path λ is larger than l_A , the stronger the turbulence, the lower thermal conductivity by electrons is. The turbulent motions, however, induces their own advective heat transport, which, for the parameters of intracluster medium (ICM) provides effective heat diffusivity that exceeds the classical Spitzer value.

Subject headings: turbulence – ISM: general – galaxies: clusters: general – MHD

1. ASTROPHYSICAL MOTIVATION

Heat transfer in turbulent magnetized plasma is an important astrophysical problem which is relevant to the wide variety of circumstances from mixing layers in the Local Bubble (see Smith & Cox 2001) and Milky way (Begelman & Fabian 1990) to cooling flows in intracluster medium (ICM) (Fabian 1994). The latter problem has been subjected to particular scrutiny as observations do not support the evidence for the cool gas (see Fabian et al. 2001). This is suggestive of the existence of heating that replenishes the energy lost via X-ray emission. Heat transfer from hot outer regions is an important process to consider in this context.

It is well known that magnetic fields can suppress thermal conduction perpendicular to their direction. The issue of heat transfer in realistic turbulent magnetic fields has been long debated (see Bakunin 2005 and references therein). An influential paper by Narayan & Medvedev (2001, henceforth NM01) obtained estimates of thermal conductivity by electrons using the Goldreich-Shidhar (1995, henceforth GS95) model of MHD turbulence with the velocity V_L at the energy injection scale L that is equal to the Alfvén velocity V_A , i.e. the turbulence with the Alfvén Mach number $M_A \equiv (V_L/V_A) = 1$. This is rather restrictive, as in the ICM $M_A > 1$ (see §4), while in other astrophysical situations $M_A < 1$. Below we discuss turbulence for both $M_A > 1$ and $M_A < 1$ and compare the heat transfer by electrons to that by turbulent fluid motions.

2. THERMAL CONDUCTIVITY: STATIC MAGNETIC FIELD

2.1. Basics of heat transfer in magnetized plasma

Following NM01, we initially disregard the dynamics of fluid motions on heat transfer, i.e. consider thermal conductivity induced by electrons moving along static magnetic fields. Magnetized turbulence in the GS95 model is anisotropic with eddies elongated along (henceforth denoted by $\parallel$) the direction of local magnetic field. Consider isotropic injection of energy at the outer scale L and dissipation at the scale $l_{\perp, \min}$, where $\perp$ denotes the direction of perpendicular to the local magnetic field. NM01 observed that the separations of magnetic field lines for $r_0 < l_{\perp, \min}$ are mostly influenced by the mo-

tions at the scale $l_{\perp, \min}$, which results in Lyapunov-type growth: $\sim r_0 \exp(l/l_{\perp, \min})$. This growth is similar to that obtained in earlier models with a single scale of turbulent motions (Rechester & Rosenbluth 1978, Chandran & Cowley 1998). This is not surprising as the largest shear that causes field line divergence is provided by the marginally damped motions at the scale around $l_{\perp, \min}$. In NM01 r_0 is associated with the size of the cloud of electrons of the electron Larmor radius $r_{Lar, electr}$. They find that the electrons should travel over the distance

$$L_{RR} \sim l_{\parallel, \min} \ln(l_{\perp, \min}/r_{Lar, electr}) \quad (1)$$

to get separated by $l_{\perp, \min}$.

Within the single-scale model which formally corresponds to $L = l_{\parallel, \min} = l_{\perp, \min}$ the scale L_{RR} is called Rechester-Rosenbluth distance. For the ICM parameters the logarithmic factor in Eq. (1) is of the order of 30, and this causes 30 times decrease of thermal conductivity for the single-scale models¹. In the multi-scale models with a limited (e.g. a few decades) inertial range the logarithmic factor stays of the same order but it does not affect the thermal conductivity, provided that $L \gg l_{\parallel, \min}$. Indeed, for the electrons to diffuse isotropically they should spread from $r_{Lar, electr}$ to L . The GS95 model of turbulence operates with field lines that are sufficiently stiff, i.e. the deviation of the field lines from their original direction is of the order unity at scale L and less for smaller scales. Therefore to get separated from the initial distance of $l_{\perp, \min}$ to a distance L (see Eq. (5) with $M_A = 1$), at which the motions get uncorrelated the electron should diffuse the distance slightly larger (as field lines are not straight) than $\sqrt{2}L$ (NM01, also see §2.3.), which is much larger than the extra travel distance $\sim 30l_{\parallel, \min}$. Explicit calculations in NM01 support this intuitive picture.

2.2. Heat Transfer for $M_A > 1$

Turbulence with $M_A > 1$ evolves along hydrodynamic isotropic Kolmogorov cascade, i.e. $V_l \sim V_L(l/L)^{1/3}$ over

¹ For the single-scale model $L_{RR} \sim 30L$ and the diffusion over distance Δ takes L_{RR}/L steps, i.e. $\Delta^2 \sim L_{RR}L$, which decreases the corresponding diffusivity coefficient $\kappa_{electr, single} \sim \Delta^2/\delta t$ by the factor of 30.

the range of scales $[L, l_A]$, where

$$l_A \approx L(V_A/V_L)^3 \equiv LM_A^{-3}, \quad (2)$$

is the scale at which the magnetic field gets dynamically important, i.e. $V_l = V_A$. This scale plays the role of the injection scale for the GS95 turbulence, i.e. $V_l \sim V_A(l_\perp/l_A)^{1/3}$, with eddies at scales less than l_A getting elongated in the direction of the local magnetic field. The corresponding anisotropy can be characterized by the relation between the semi-major axes of the eddies

$$l_\parallel \sim L(l_\perp/L)^{2/3}M_A^{-1}, \quad M_A > 1, \quad (3)$$

where $\parallel$ and $\perp$ are related to the direction of the local magnetic field. In other words, for $M_A > 1$, the turbulence is still isotropic at the scales larger to l_A , but develops $(l_\perp/l_A)^{1/3}$ anisotropy for $l < l_A$.

For electron mean free path $\lambda \gg l_A$, electrons stream freely over the distance of l_A . For electrons at distance $l_{\perp, \min}$ to get separated by L the required travel is the random walk with the step l_A , i.e. the mean-squared displacement of a thermal electron till it enters an independent large-scale eddy $\Delta^2 \sim l_A^2(L/l_A)$, where L/l_A is the number of steps. These steps require time $\delta t \sim (L/l_A)l_A/C_1v_{electr}$, where v_{electr} is electron thermal velocity and the coefficient $C_1 = 1/3$ accounts for 1D character of motion along magnetic field lines. Thus the electron diffusivity coefficient is

$$\kappa_{electr} \equiv \Delta^2/\delta t \approx (1/3)l_A v_{electr}, \quad l_A < \lambda, \quad (4)$$

which for $l_A \ll \lambda$ constitutes a substantial reduction of conductivity compared to its Spitzer (unmagnetized) value $\kappa_{spitzer} = \lambda v_{electr}$. We assumed in Eq. (4) that $L \gg 30l_{\parallel, \min}$ (see §2.1).

For $\lambda \ll l_A \ll L$, $\kappa_{electr} \approx 1/3\kappa_{spitzer}$ as both the L_{RR} and the additional distance for electron to diffuse because of magnetic field being stiff at scales less than l_A are negligible compared to L . For $l_A \rightarrow L$, when magnetic field has rigidity up to the scale L , it gets around 1/5 of the Spitzer value according to NM01.

Note, that even dynamically unimportant magnetic fields do influence heat conductivity over short time intervals. For instance, over time interval less than $l_A^2/C_1\kappa_{spitzer}$ the diffusion happens along stiff magnetic field lines and the difference between parallel and perpendicular diffusivities is large². This allows the transient existence of sharp small-scale temperature gradients.

2.3. Heat Transfer for $M_A < 1$

It is intuitively clear that for $M_A < 1$ turbulence should be anisotropic from the injection scale L . In fact, at large scales the turbulence is expected to be *weak*³ (see Lazarian & Vishniac 1999, henceforth LV99). Weak turbulence is characterized by wavepackets that

² The relation between the mean squared displacements perpendicular to magnetic field $\langle y^2 \rangle$ and the displacements x along magnetic field for $x < l_A$ can be obtained through the diffusion equation approach in §2.3 and Eq. (3). This gives $\langle y^2 \rangle^{1/2} \sim \frac{x^{3/2}}{3^{3/2}L^{1/2}}M_A^3$.

³ The terms “weak” and “strong” turbulence are accepted in the literature, but can be confusing. As we discuss later at smaller scales at which the turbulent velocities decrease the turbulence becomes *strong*. The formal theory of weak turbulence is given in Galtier et al. (2000).

do not change their $l_\parallel$, but develop structures perpendicular to magnetic field, i.e. decrease $l_\perp$. This cannot proceed indefinitely, however. At some small scale the GS95 condition of *critical balance*, i.e. $l_\parallel/V_A \approx l_\perp/V_l$, becomes satisfied. This perpendicular scale l_{trans} can be obtained substituting the scaling of weak turbulence (see LV99) $V_l \sim V_L(l_\perp/L)^{1/2}$ into the critical balance condition. This provides $l_{trans} \sim LM_A^2$ and the corresponding velocity $V_{trans} \sim V_L M_A$. For scales less than l_{trans} the turbulence is *strong* and it follows the scalings of the GS95-type, i.e. $V_l \sim V_L(L/l_\perp)^{-1/3}M_A^{1/3}$ and

$$l_\parallel \sim L(l_\perp/L)^{2/3}M_A^{-4/3}, \quad M_A < 1. \quad (5)$$

For $M_A < 1$, magnetic field wandering in the direction perpendicular to the mean magnetic field (along y-axis) can be described by $d\langle y^2 \rangle/dx \sim \langle y^2 \rangle/l_\parallel$ (LV99), where⁴ $l_\parallel$ is expressed by Eq. (5) and one can associate $l_\perp$ with $2\langle y^2 \rangle$

$$\langle y^2 \rangle^{1/2} \sim \frac{x^{3/2}}{3^{3/2}L^{1/2}}M_A^2, \quad l_\perp < l_{trans} \quad (6)$$

For weak turbulence $d\langle y^2 \rangle/dx \sim LM_A^4$ (LV99) and thus

$$\langle y^2 \rangle^{1/2} \sim L^{1/2}x^{1/2}M_A^2, \quad l_\perp > l_{trans}. \quad (7)$$

Eq. (6) differs by the factor M_A^2 from that in NM01, which reflects the gradual suppression of thermal conductivity perpendicular to the mean magnetic field as the magnetic field gets stronger. Physically this means that for $M_A < 1$ the magnetic field fluctuates around the well-defined mean direction. Therefore the thermal conduction gets anisotropic with the coefficient of thermal conduction parallel to the mean field $\kappa_{\parallel, electr} \approx 1/3\kappa_{spitzer}$ being larger than $\kappa_{\perp, electr}$ for the thermal conductivity in the perpendicular direction.

Consider the coefficient $\kappa_{\perp, electr}$ for $M_A \ll 1$. As NM01 showed, electrons become uncorrelated if they are displaced over the distance L in the direction perpendicular to magnetic field. To do this, an electron has first to travel L_{RR} (see Eq. (1)), where Eq. (5) relates $l_{\parallel, \min}$ and $l_{\perp, \min}$. Similar to the case in §2.1, for $L \gg 30l_{\parallel, \min}$, the additional travel arising from the logarithmic factor is negligible compared to the overall diffusion distance L . At larger scales electron has to diffuse $\sim L$ in the direction parallel to magnetic field to cover the distance of LM_A^2 in the direction perpendicular to magnetic field direction. Therefore the separation of electrons over the turbulence driving scale L perpendicular to the magnetic field direction requires $L/LM_A^2 = M_A^{-2}$ random steps. If $\lambda \ll L$ the diffusion over L requires time $\delta t \sim L^2/(M_A^2 D_\parallel)$, where $D_\parallel$ is the diffusion coefficient which is $v_{electr}\lambda/3$. As a result

$$\kappa_{\perp, electr} \equiv L^2/\delta t \approx 1/3v_{electr}\lambda M_A^2, \quad M_A < 1, \quad (8)$$

where we disregarded the distance to travel in the direction perpendicular mean magnetic field, i.e. L , compared to the distance to travel parallel to magnetic field, i.e. LM_A^2 . For M_A of the order of unity this is not accurate and one should account for the actual 3D displacement (see NM01 and §2.1).

⁴ The fact that one gets $l_{\parallel, \min}$ in Eq. (1) is related to the presence of this scale in this diffusion equation.




FIG. 1.— Sonic Mach number M_s is plotted against the Alfvén Mach number M_A . The heat transport is dominated by the dynamics of turbulent eddies is above the curve and by thermal conductivity of electrons is below the curve. Here λ is the mean free path of the electron, L is the driving scale, and $\alpha = (m_e/m_p)^{1/2}$, $\beta \approx 4$. The panel in the right upper corner of the figure illustrates heat transport for the parameters for a cool core Hydra cluster using data from EV06 (point “CC”), “I” corresponds to the illustrative model in EVP05.

3. FLUID VERSUS ELECTRON MOTIONS

Turbulent motions themselves can advectively transport heat. In Cho et al. (2003) we dealt with the turbulence with $M_A \sim 1$ and estimated

$$\kappa_{dynamic} \approx C_{dyn} L V_L, \quad M_A > 1, \quad (9)$$

where $C_{dyn} \sim 0(1)$ is a constant, which for hydro turbulence is around $1/3$ (Lesieur 1990). For fully ionized non-degenerate plasma we assume $C_{dyn} \approx 2/3$ to account for the advective heat transport by both protons and electrons⁵. Thus eq. (9) covers the cases of both $M_A > 1$ up to $M_A \sim 1$. For $M_A < 1$ one can estimate $\kappa_{dynamic} \sim d^2 \omega$, where d is the random walk of the field line over the wave period $\sim \omega^{-1}$. As the weak turbulence at scale L evolves over time $\tau \sim M_A^{-2} \omega^{-1}$, $\langle y^2 \rangle$ is the result of the random walk with a step d , i.e. $\langle y^2 \rangle \sim (\tau \omega) d^2$. According to eq.(6) and (7), the field line is displaced over time τ by $\langle y^2 \rangle \sim L M_A^4 V_A \tau$. Combining the two one gets $d^2 \sim L M_A^3 V_L \omega^{-1}$, which provides $\kappa_{dynamic}^{weak} \approx C_{dyn} L V_L M_A^3$, which is similar to the diffusivity arising from strong turbulence at scales less than l_{trans} , i.e. $\kappa_{dynamic}^{strong} \approx C_{dyn} l_{trans} V_{trans}$. The total diffusivity is the sum of the two, i.e. for plasma

$$\kappa_{dynamic} \approx (\beta/3) L V_L M_A^3, \quad M_A < 1, \quad (10)$$

where $\beta \approx 4$.

The schematic of the parameter space for $\kappa_{electr} < \kappa_{dynamic}$ is shown in Fig 1, where the the Mach number M_s and the Alfvén Mach number M_A are the variables. For $M_A < 1$, the ratio of thermal conductivities arising from fluid and electron motions is $\kappa_{dynamic}/\kappa_{electr} \sim \beta \alpha M_s M_A (L/\lambda)$ (see Eqs. (8) and (10)), the square root of the ratio of the electron to proton mass $\alpha = (m_e/m_p)^{1/2}$, which provides the separation line between the two regions in Fig. 1, $\beta \alpha M_s \sim$

$(\lambda/L) M_A^{-1}$. For $1 < M_A < (L/\lambda)^{1/3}$ the mean free path is less than l_A which results in κ_{electr} being some fraction of $\kappa_{spitzer}$, while $\kappa_{dynamic}$ is given by Eq. (9). Thus $\kappa_{dynamic}/\kappa_{electr} \sim \beta \alpha M_s (L/\lambda)$, i.e. the ratio does not depend on M_A (horizontal line in Fig. 1). When $M_A > (L/\lambda)^{1/3}$ the mean free path of electrons is constrained by l_A . In this case $\kappa_{dynamic}/\kappa_{electr} \sim \beta \alpha M_s M_A^3$ (see Eqs. (9) and (4)). This results in the separation line $\beta \alpha M_s \sim M_A^{-3}$ in Fig. 1.

4. TURBULENCE AND HEAT TRANSFER IN ICM

It is generally believed that ICM is turbulent. The considerations below can be used as guidance. In unmagnetized plasma with the ICM temperatures $T \sim 10^8$ K and density 10^{-3} cm^{-3} the diffusivity $\nu_{B=0} \sim v_{ion} \lambda_{ion}$, where v_{ion} and λ_{ion} are the velocity of an ion and its mean free path, respectively, would make the Reynolds number $Re \equiv L V_L / \nu$ of the order of 30. This is barely enough for the onset of turbulence. For the sake of simplicity we assume that ion mean free path coincides with the proton mean free path and both scale as $\lambda \approx 3 T_3^2 n_{-3}^{-1} \text{ kpc}$, where the temperature $T_3 \equiv kT/3 \text{ keV}$ and $n_{-3} \equiv n/10^{-3} \text{ cm}^{-3}$. This provides λ of the order of $0.8\text{--}1 \text{ kpc}$ for the ICM (see NM01).

It is accepted, however, that magnetic fields decrease the diffusivity. Somewhat naively assuming the maximal scattering rate of an ion, i.e. scattering every orbit (the so-called Bohm diffusion limit) one gets the viscosity perpendicular to magnetic field $\nu_{\perp} \sim v_{ion} r_{Lar,ion}$, which is much smaller than $\nu_{B=0}$, provided that the ion Larmor radius $r_{Lar,ion} \ll \lambda_{ion}$. For the parameters of the ICM this allows essentially inviscid motions⁶ of magnetic lines parallel to each other, e.g. Alfvén motions.

In spite of the substantial progress in understading of the ICM (see Enßlin, Vogt & Pfrommer 2005, henceforth EVP05, Enßlin & Vogt 2006, henceforth EV06 and references therein), the basic parameters of ICM turbulence are known within the factor of 3 at best. For instance, the estimates of injection velocity V_L varies in the literature from 300 km/s to 10^3 km/s , while the injection scale L varies from 20 kpc to 200 kpc , depending whether the injection of energy by galaxy mergers or galaxy wakes is considered. EVP05 considers an *illustrative* model in which the magnetic field with the $10 \mu\text{G}$ fills 10% of the volume, while 90% of the volume is filled with the field of $B \sim 1 \mu\text{G}$. Using the latter number and assuming $V_L = 10^3 \text{ km/s}$, $L = 100 \text{ kpc}$, and the density of the hot ICM is 10^{-3} cm^{-3} , one gets $V_A \approx 70 \text{ km/s}$, i.e. $M_A > 1$. Using the numbers above, one gets $l_A \approx 30 \text{ pc}$ for the 90% of the volume of the hot ICM, which is much less than λ_{ion} . The diffusivity of ICM plasma gets $\nu = v_{ion} l_A$ which for the parameters above provides $Re \sim 2 \times 10^3$, which is enough for driving superAlfvénic turbulence at the outer scale L . However, as l_A increases as $\propto B^3$, Re gets around 50 for the field of $4 \mu\text{G}$, which is at the

⁵ This gets clear if one uses the heat flux equation $q = -\kappa_c \nabla T$, where $\kappa_c = n k_B \kappa_{dynamic/electr}$, n is electron number density, and k_B is the Boltzmann constant, for both electron and advective heat transport.

⁶ A regular magnetic field $B_{\lambda} \approx (2mkT)^{1/2} c/(e\lambda)$ that makes $r_{Lar,ion}$ less than λ and therefore $\nu_{\perp} < \nu_{B=0}$ is just 10^{-20} G . Turbulent magnetic field with many reversals over $r_{Lar,ion}$ does not interact efficiently with a proton, however. As the result, the protons are not constrained until l_A gets of the order of $r_{Lar,ion}$. This happens when the turbulent magnetic field is of the order of $2 \times 10^{-9} (V_L/10^3 \text{ km/s}) \text{ G}$. At this point, the step for the random walk is $\sim 2 \times 10^{-6} \text{ pc}$ and the Reynolds number is 5×10^{10} .

border line of exciting turbulence⁷. However, the regions with higher magnetic fields (e.g. 10 μG) can support Alfvénic-type turbulence with the injection scale l_A and the injection velocities resulting from large-scale shear $V_L(l_A/L) \sim V_L M_A^{-3}$.

For the regions of $B \sim 1 \mu\text{G}$ the value of l_A is smaller than the mean free path of electrons λ . According to Eq. (4) the value of κ_{electr} is 100 times smaller than $\kappa_{spitzer}$. On the contrary, $\kappa_{dynamic}$ for the ICM parameters adopted will be $\sim 30\kappa_{spitzer}$, which makes the dynamic diffusivity the dominant process. This agrees well with the observations in Voigt & Fabian (2004). Fig. 1 shows the dominance of advective heat transfer for the parameters of the cool core of Hydra A ($B = 6 \mu\text{G}$, $n = 0.056 \text{ cm}^{-3}$, $L = 40 \text{ kpc}$, $T = 2.7 \text{ keV}$ according to EV06), point “CC”, and for the illustrative model in EVP05, point “T”, for which $B = 1 \mu\text{G}$.

Note that our stationary model of MHD turbulence in §2 is not directly applicable to transient wakes behind galaxies. The ratio of the damping times of the hydro turbulence and the time of straightening of the magnetic field lines is $\sim M_A^{-1}$. Thus, for $M_A > 1$, the magnetic field at scales larger than l_A will be straightening gradually after the hydro turbulence has faded away over time L/V_L . The process can be characterized as injection of turbulence at velocity V_A but at scales that increase linearly with time, i.e. as $l_A + V_A t$. The study of heat transfer in transient turbulence and magnetic field “regularly” stretched by passing galaxies will be provided elsewhere.

5. DISCUSSION AND SUMMARY

In the paper above we attempted to describe the heat transfer by electron and turbulent motions for $M_A < 1$ and $M_A > 1$. Unlike earlier papers, we find that turbulence may both enhance heat conduction and suppress it. For instance, when λ gets larger than l_A the conductivity of the medium $\sim M_A^{-3}$ and therefore the turbulence *inhibits* heat transfer, provided that $\kappa_{electr} > \kappa_{dynamic}$. Along with the plasma effects that we mention below, this effect can, indeed, support sharp temperature gradients in hot plasmas with weak magnetic field.

As discussed above, rarefied plasma, e.g. ICM plasma, has large viscosity for motions parallel to magnetic field

and marginal viscosity for motions that induce perpendicular mixing. Thus fast dissipation of sound waves in the ICM does not contradict the medium being turbulent. The later may be important for the heating of central regions of clusters caused by the AGN feedback (see Churazov et al. 2001, Nusser, Silk & Babul 2006 and more references in EV06). Note, that models that include both heat transfer from the outer hot regions and an additional heating from the AGN feedback look rather promising (see Ruszkowski & Begelman 2002, Piffaretti & Kaastra 2006). We predict that the viscosity for 1 μG regions is less than for 10 μG regions and therefore heating by sound waves (see Fabian et al. 2005) could be more efficient for the latter. Note, that the plasma instabilities in collisionless magnetized ICM arising from compressive motions (see Schekochihin & Cowley 2006, Lazarian & Beresnyak 2006) can resonantly scatter electrons and protons and decrease λ for both species compared to the classical plasma values (λ gets different for electrons and protons in this case). This decreases further κ_{electr} compared to $\kappa_{spitzer}$ but increases Re . In addition, we disregarded mirror effects that can reflect electrons back (see Mal'ushkin & Kulsrud 2001 and references therein), which can further decrease κ_{electr} .

All in all, we have shown that it is impossible to characterize the heat transfer of magnetized plasma by a single fraction of Spitzer's value. The actual heat transport depends on sonic and Alfvén Mach numbers of turbulence and may be much higher and much lower than the classical one. As the result, turbulence can inhibit or enhance heat conductivity depending on the plasma magnetization and turbulence driving. Our study indicates that in many cases related to ICM the advective heat transport by dynamic turbulent eddies dominates thermal conductivity.

Acknowledgments. The work is supported by the NSF grant AST 0307869 and the NSF Center for Magnetic Self-Organization in Laboratory and Astrophysical Plasmas. I thank Ethan Vishniac for discussions of diffusion in the weak turbulence case, Pat Diamond for insight into plasma effects, Torsten Enßlin and Andrey Beresnyak for helpful comments.

down the turbulence.

REFERENCES

- Bakunin, O.G. 2005, Plasma Phys. and Contr. Fus. 47, 1857
 Begelman, M. & Fabian, A. 1990, MNRAS, 244, 26
 Chandran, B. & Cowley, S. 1998, Phys. Rev. Lett., 80, 3077
 Cho, J. & Lazarian, A. 2002, Phys. Rev. Lett., 88, 5001
 Cho, J., Lazarian, A., Honein, A., Kassions, S., & Moin, P. 2003, ApJ, 589, L77
 Churazov, E., Bruggen, M., Kaiser, C., Bohringer, H., & Forman W. 2001, ApJ, 554, 261
 Galtier, S., Nazarenko, S., Newel, A. & Pouquet, A. 2000, J. Plasma Phys., 63, 447
 Enßlin, T., Vogt, C. & Pfrommer, C. 2005, in The Magnetized Plasma in Galaxy Evolution, Eds. K.T. Chyzy, K. Otminowska-Mazur, M. Soida and R.-J. Dettmar, Kracow, p. 231
 Enßlin, T., & Vogt, C. 2006, ApJ, in press
 Fabian, A.C. 1994, ARA&A, 32, 277
 Fabian, A.C., Mushotzky, R.F., Nulsen, P.E.J., & Peterson, J.R. 2001, MNRAS, 321, L20
 Fabian, A.C., Reynolds, C.S., Taylor, G.B. & Dunn, R.J. 2005, MNRAS, 363, 891
 Goldreich, P. & Sridhar, S. 1995, ApJ, 438, 763
 Lazarian, A. & Beresnyak, A. 2006, MNRAS, submitted
 Lazarian, A. & Vishniac, E. 1999, ApJ, 517, 700
 Mal'ushkin, L. & Kulsrud, R. 2001, ApJ, 549, 402
 Lesieur, M. 1990, *Turbulence in fluids : stochastic and numerical modelling*, 2nd. rev. ed. (Dordrecht; Kluwer)
 Narayan, R., & Medvedev M. 2001, ApJ, 562, L129
 Nusser, A., Silk, J. & Babul, A. 2006, astro-ph/0602566
 Piffaretti, R. & Kaastra, J. 2006, astro-ph/0602376
 Rechester, A., & Rosenbluth, M. 1978, Phys. Rev. Lett., 40, 38
 Ruszkowski, M. & Begelman, M.C. 2002, ApJ, 581, 223
 Schekochihin, A. & Cowley, S. 2006, astro-ph/0601246
 Smith, R. & Cox, D. 2001, ApJS, 134, 283
 Voigt, L.M. & Fabian, A.C. 2004, MNRAS, 347, 1130